

Research Use Only. Not for use in diagnostic procedures.

Bioware[®] Brite Cell Line HT1080 Red-FLuc

Product No.: BW 128092

Material Provided

Cells: 2 x 1 mL frozen aliquots (BW 128092V)

Format: 1.0 x 10⁶ cells / mL in 95% FBS, 5% DMSO

DESIGNATION	HT1080 Red-FLuc
Tissue	Human fibrosarcoma
Source of Parental Line	ATCC (CCL-121)
Gene Transfer Vehicle	Red-FLuc-Puro 3d generation lentivirus
Bioluminescence In Vitro	At least 30,000 photons/cell/sec. Exact number will vary depending on imaging and culturing conditions.
Recommended Media and FBS	Eagle's MEM ATCC Cat. No. 30-2007. Supplement the above with 10% Hyclone Fetal Bovine Serum (FBS) GE HealthCare Cat. No. SH300071
Recommended Storage Conditions	Remove frozen cells from dry ice packaging and immediately place cells at a temperature below -130° C, preferably in liquid nitrogen vapor, until ready to use.
Average Doubling Time	22 hours
Other Recommendations	When initially thawing, use T25 flask or 10cm plate. Cells should be ready to expand within 1-4 days. Antibiotics can be used in the media if desired after the initial thaw. (puromycin at 2ug/mL). Refer to Cell Culture Guidelines for more detailed instructions.

PerkinElmer, Inc.
940 Winter Street
Waltham, MA 02451 USA
P: (800) 762-4000 or
(+1) 203-925-4602
www.perkinelmer.com



For a complete listing of our products, visit www.perkinelmer.com.

Copyright ©2015, PerkinElmer, Inc. All rights reserved. PerkinElmer[®] is a registered trademark of PerkinElmer, Inc. All other trademarks are the property of their respective owners.

The Features

Perkin Elmer Bioware® Brite cell line models offer researchers the ability to:

- Monitor early tumor development
- Monitor tumor growth and metastases *in vivo*
- Quantify tumor burden in the whole animal
- Follow responses to therapeutic treatments non-invasively in longitudinal studies using the same cohorts of mice

Murine Pathogen Free

All Perkin Elmer cell lines are confirmed to be pathogen free by the IMPACT Profile I (PCR) at the University of Missouri Research Animal Diagnostic and Investigative Laboratory.

Cell Line Stability

Cell may undergo genotypic changes resulting in reduced responsiveness over time in normal cell culture conditions. Genetic instability is a biological phenomenon that occurs in all stably transfected cells. Therefore, it is recommended to prepare an adequate number of frozen stock at early passages.

Product Warranty

PerkinElmer warrants that cells will be viable upon shipment from PerkinElmer for a period of thirty days, provided they have been properly stored and handled during this period.

Human Fibrosarcoma Cancer Cell Line: HT1080 Red-FLuc

HT1080 Red-FLuc is a luciferase expressing cell line which was stably transfected with firefly luciferase gene from *Luciola Italica* (Red-FLuc). The cell line was established by transducing lentivirus containing Red-FLuc luciferase under the control of human ubiquitin C promoter. These cells will serve as a new tool to detect drug efficacy *in vitro* and *in vivo* with high sensitivity.



Bioluminescence image
of HT1080 Red-FLuc
subcutaneous tumor

Growth Curve of HT1080 Red-FLuc Cells




Figure 1. 5×10^4 cells were plated on 6cm plate and the total numbers of cells were counted every 24 h using a Nexcelom automatic cell counter.

In Vitro BLI Signal Stability





Figure 2. 5×10^4 cells were plated per well in 24-well plates. Cells were incubated at 37 °C for recovery overnight and luciferase assay was performed using the PerkinElmer IVIS[®] SpectrumCT. Each experiment was done in quadruplicates. The cells were maintained in continuous culture over four weeks and weekly luciferase assay was performed. Bioluminescence data was analyzed using the Living Image 4.0 software.

Subcutaneous Tumor Growth in a Nu/nu Mouse




Figure 3. 1×10^6 HT1080-Red-FLuc cells were injected subcutaneously into the dorsal region near the thigh of female nu/nu mouse. Tumor growth was monitored for luciferase expression using the PerkinElmer IVIS[®] Spectrum at various time points. Mice were imaged 10 minutes post i.p. injection of luciferin at 150mg/kg at various time points. The image above shows tumor growth from a representative mouse.

For more information on our *in vivo* imaging agents, please visit our website: www.perkinelmer.com/bioware.

This product is sold for *in vivo* animal research use only and is not intended for any diagnostic use or procedures. Excluding purchases by authorized PerkinElmer distributors, this product is sold for use by the original purchaser and is not for resale. Unless otherwise agreed to in writing by PerkinElmer pursuant to a separate written agreement, no commercial use of this product is allowed. “Commercial use” means any and all uses of this product and/or its derivatives by a party for money or other consideration and may include without limitation: (1) product manufacture; (2) providing services, information or data to another party for remuneration; and/or (3) resale of the product or its derivatives, whether or not such product or derivatives are resold for use in research. Commercial use does not include the original purchaser providing the product to its contractor solely for use on the original purchaser’s research; provided that all product materials are returned to the original purchaser and/or destroyed by the contractor upon completion of such project.